

Australian and New Zealand College of Anaesthetists (ANZCA)

Guidelines for the Safe Management and Use of Medications in Anaesthesia

Background Paper

1. PURPOSE OF REVIEW

PS51 *Guidelines for the Safe Administration of Injectable Drugs in Anaesthesia* was promulgated in 2009 and is now due for review. PS51 is referenced and referred to by healthcare facility administration, fellows, and trainees. The current revision has aimed to assist administrators and practitioners in producing the safest environment for patients and staff, as well as accommodate the potentially conflicting needs for ready availability of medications and the opportunity for misuse.

2. BACKGROUND

As a result of a number of lethal outcomes from the diversion and misuse of propofol, hospitals and jurisdictions have introduced different solutions to a common problem. Some solutions have included restricting access to propofol. As propofol is a drug used for the induction and maintenance of anaesthesia it must be immediately available and thus restricting access has the potential to impact on the safe conduct of anaesthesia. However, propofol should be securely stored at all times when not immediately required, for example overnight and between operating sessions.

The accompanying professional document provides guidance to the profession and hospitals to avoid the implementation of a multitude of solutions across Australian states and NZ. In addition, it makes recommendations aimed at reducing drug errors and risks of cross-contamination between patients.

Any change to regulatory requirements in light of recent reviews of various drugs, poisons and controlled substances legislation will also need to be taken into account.

It is recognised that there are jurisdictional differences between Australia and New Zealand, particularly with respect to purchasing authority, labelling of medications and standards pertaining to user applied labels.

3. DISCUSSION

In reviewing the accompanying guidelines, the following issues were considered:

- 3.1 **Title of the document** - The title was modified to reflect the broader intent of the guidelines beyond simply the administration of medications and instead to include safety in management and use of medications.
- 3.2 **Drug errors** - In recognition of the range of reported drug errors^{1,2}, which may reflect the tip of the iceberg, attempts have been made to address each potential factor. These have included:



- 3.2.1 Identification of vials and ampoules – with the similarity of some presentations more than just a cursory check of labels is required. Notwithstanding this, it was agreed that there should be standardisation of labelling³ for all classes of drugs by manufacturers. There is already a standard for syringe labelling and it makes sense to apply the same system to ampoules and vials. Special attention should be directed towards standardisation of presentation of the information with regard to font, size, and colour of print.
- 3.2.2 Labelling of syringes by applying labels compliant with the relevant National Standard⁴. In the specific situation where a medication is drawn up and immediately administered as a bolus, without leaving the hands of the practitioner that prepared it, labelling is not required. In all other cases labelling is required as per the National Standard for User Applied Labelling in Australia. ANZCA has released a joint statement with the ACSQHC supporting user-applied labelling standardisation for all injectable medicines and fluids⁴.
In New Zealand PHARMAC is considering labelling recommendations.
- 3.2.3 With the increasing use of infusions for both anaesthesia as well as for regional analgesia there is the potential for drug errors to occur. To minimise these errors, it is advisable to ensure that devices delivering intravenous medications are readily differentiated from those delivering local anaesthetics. Colour coding for different routes and compliance with Labelling Standards is required in Australia. With the coming introduction of the new small bore neuraxial standard the likelihood of such misconnections should be diminished. Infusion devices with Dosage Error Reduction Software may offer a safety benefit⁵.
- 3.2.4 Patient identification and “open” versus “closed” practice environments.

Anaesthetists must be aware of the difference between an open and closed practice environment as there are implications for ensuring delivery of the correct medication to the correct patient.

In a closed-practice environment:

- the identity of the patient is beyond doubt – that is, when a single patient is receiving an injectable medicine, there is no possibility that the patient’s identity is unknown, and the medicine is prepared in the presence of the patient
- the identities of members of the patient care team are recorded
- the medicine is administered completely within the closed-practice environment. In this case, patient identifiers are not required and a pre-printed abbreviated label may be used.

Examples of closed-practice environments are operating rooms, endoscopy rooms, catheterisation laboratories and radiology suites.

In an open-practice environment there are multiple patients in the same area and full patient and user identification is required for user-applied labelling. The PACU is an open environment as there are numerous possible candidates for the drug drawn up and to be administered⁶.

- 3.2.5 Purchasing decisions on anaesthesia drugs should be made in consultation with a designated safety officer in the department of anaesthesia or other nominated individual⁷. Purchase choices should be made in accordance with the principles articulated in PS51, relevant jurisdictional requirements, and with regard to local practice environments.

3.3 **Infection control** – the potential for cross-contamination and the need for maintaining sterility of drugs drawn up into syringes can be addressed by adherence to the principles of aseptic non touch technique⁸. The maintenance of critical micro aseptic fields is facilitated by capping of syringes.

3.4 **Storage of drugs** – While the storage and documentation of medications must comply with the

regulations of the local jurisdictional authority, the anaesthetist must be able to directly and independently access sufficient quantities of critical drugs. Such drugs would include propofol, adrenaline, muscle relaxants, and vasopressors.

- 3.5 Interruptions to supply** –Recently, availability of a number of medications has been restricted due to problems with manufacture and supply, or concerns about potential contamination. Development of an “essential medicines list” has been proposed but cannot be mandated. The existence of dual pharmaceutical suppliers of a particular medication assists in minimising disruption to availability.
- 3.6 Automated medication dispensing systems** –many healthcare facilities are using medication and storage systems that allow tracking of stock levels, access identification and patient prescription. They usually require fingerprint or other identification to access and thus whilst providing enhanced security, they are not suitable when medications are required immediately. It is strongly recommended that a supply of medications required in an emergency or urgently is readily available outside an automated storage device.
- 3.7 Volatile Agents** - Volatile agents have been identified as drugs of abuse⁹ and thus must be stored securely. Case reports of substitution of a cleaning solution for a volatile agent indicate the importance of appropriate storage of partially filled bottles and appropriate disposal of empty containers¹⁰.

4. SUMMARY

The current revision has considered the changing environment since the promulgation of the previous version and has reviewed the issues relevant to the broader aspects of the safe management and use of medications in anaesthesia. The recommendations in the accompanying guidelines should assist healthcare facilities, administrators, and practitioners to provide safer conditions for their patients.

REFERENCES

1. Nanji KC, Patel A, Shaikh S, Seger DL, Bates DW. [Evaluation of perioperative medication errors and adverse drug events](#). Anesthesiology. 2016;124(1): 25-34. doi: 10.1097/ALN.0000000000000904
2. Webster CS, Merry AF, Larsson L, McGrath KA, Weller J. The frequency and nature of drug administration error during anaesthesia. Anaesth Intensive Care [serial online]. 2001;29(5):494-500. From: <http://www.aaic.net.au.ezproxy.anzca.edu.au/Document/?D=2000210>. Accessed 12 October 2016
3. Department of Health, Therapeutic Goods Administration [Internet] 2016 From: <https://www.legislation.gov.au/Details/F2016L01285>
4. Australian Commission on Safety and Quality in Health Care. A joint statement supporting user-applied labelling standardization for all injectable medicines and fluids by the Australian Commission on Safety and Quality in Health Care, and the Australian and New Zealand College of Anaesthetists [Internet]. 2017. From <https://www.safetyandquality.gov.au/wp-content/uploads/2017/01/Joint-statement-supporting-user-applied-labelling-standardisation-for-all-injectable-medicines-and-fluids-January-2017.pdf>
5. Pang RKY, Kong DCM, deClifford J-M, Lam SS, Leung BK. [Smart Infusion Pumps Reduce Intravenous Medication Administration Errors at an Australian Teaching Hospital](#) Journal of Pharmacy Practice and Research. 2011; 41: 192-5.
6. Australian Commission on Safety and Quality in Health Care. National standard for user-applied labelling of injectable medicines, fluids and lines [Internet]. 2016. From: <https://www.safetyandquality.gov.au/wp-content/uploads/2015/09/National-Standard-for-User-Applied-Labelling-Aug-2015.pdf>. Accessed 12 October 2016.

7. Australian Commission on Safety and Quality in Health Care. Safety and Quality Improvement Guide Standard 4: Medication Safety [Internet] 2012 From: https://www.safetyandquality.gov.au/wp-content/uploads/2012/10/Standard4_Oct_2012_WEB.pdf Accessed 26 October 2016.
8. National Health and Medical Research Council [Internet] 2010 From: <https://www.nhmrc.gov.au/book/australian-guidelines-prevention-and-control-infection-healthcare-2010/b1-7-2-core-infection-co>
9. Wilson JE, Kiselanova N, Stevens Q, Lutz R, Mandler T, Tran ZV, Wischmeyer PE. [A survey of inhalational anaesthetic abuse in anaesthesia training programmes](#). Anaesthesia 2008; 63: 616-670
10. Fry R, Wilton N, Boyes G. Anaesthesia and Intensive Care 2015;43(3):419-20

DOCUMENT DEVELOPMENT GROUP

Dr Phillipa Hore, FANZCA, Chair Safety and Quality Committee, Vic.

Dr Peter Roessler, FANZCA, Director of Professional Affairs (Professional Documents), Vic

Prof David A Scott, FANZCA, FFPMANZCA. President ANZCA

Dr David Bramley, FANZCA

Dr Rob Fry, FANZCA

In addition, the following were consulted:

Safety and Quality Committee

ANZCA regional and national committees

Australian Society of Anaesthetists

Faculty of Pain Medicine Board and regional committees

ANZCA Trainee Committee

ANZCA Special Interest Groups (SIGs)

Professional documents of the Australian and New Zealand College of Anaesthetists (ANZCA) are intended to apply wherever anaesthesia is administered and perioperative medicine practised within Australia and New Zealand. It is the responsibility of each practitioner to have express regard to the particular circumstances of each case, and the application of these ANZCA documents in each case. It is recognised that there may be exceptional situations (for example, some emergencies) in which the interests of patients override the requirement for compliance with some or all of these ANZCA documents. Each document is prepared in the context of the entire body of the College's professional documents, and should be interpreted in this way.

ANZCA professional documents are reviewed from time to time, and it is the responsibility of each practitioner to ensure that he or she has obtained the current version which is available from the College website (www.anzca.edu.au). The professional documents have been prepared having regard to the information available at the time of their preparation, and

practitioners should therefore take into account any information that may have been published or has become available subsequently.

Whilst ANZCA endeavours to ensure that its professional documents are as current as possible at the time of their preparation, it takes no responsibility for matters arising from changed circumstances or information or material which may have become available subsequently.

Promulgated: 2017
Date of current document: April 2017

© Copyright 2017 – Australian and New Zealand College of Anaesthetists. All rights reserved.

This work is copyright. Apart from any use as permitted under the Copyright Act 1968, no part may be reproduced by any process without prior written permission from ANZCA. Requests and inquiries concerning reproduction and rights should be addressed to the Chief Executive Officer, Australian and New Zealand College of Anaesthetists, 630 St Kilda Road, Melbourne, Victoria 3004, Australia. Website: www.anzca.edu.au email: ceoanzca@anzca.edu.au

ANZCA website: www.anzca.edu.au