



Review Resource Document 18 (2011)

WELFARE OF ANAESTHETISTS SPECIAL INTEREST GROUP

LATEX ALLERGY

An Occupational Hazard for Anaesthetists

INTRODUCTION

Natural rubber latex is the milky sap from the rubber tree, *Hevea brasiliensis*.

The protein portion is responsible for the type I hypersensitivity: urticaria which can lead to anaphylaxis.

Chemicals give latex its elasticity, and these cause the type 4 (cell mediated) contact dermatitis.

ASSOCIATED RISK FACTORS FOR LATEX ALLERGY

- Atopy
- Exposure to Latex
 - Latex gloves are preferred for strength and exclusion of viral particles. Some synthetic gloves, eg vinyl, are not as strong or impermeable, and should not be used as protection for blood and body fluids.
- Associated food allergies:
 - Banana, avocado, chestnut, kiwi fruit, tomatoes, potatoes, stone fruits.
 - Reactions to foods may precede recognition of latex allergy and also the reverse.

WHY ARE POWDERED LATEX GLOVES A PROBLEM?

Latex Aeroallergens

Latex allergens are absorbed onto the cornstarch powder, and then become a source of airborne latex allergen, known as latex aeroallergens.

Latex aeroallergen sensitises individuals working in this environment more effectively than wearing latex gloves.

ANAESTHETISTS WITH LATEX ALLERGY

- Seek good quality occupational advice in view of the high risk work environment.
- There is no treatment available at present apart from latex avoidance, which must include avoiding the inhalation of powder from latex gloves.



- The best hope for continued anaesthetic practice is to seek early diagnosis.
- The severity of the reaction to latex must be determined.
 - Those with severe symptoms may not be able to work in acute health care environment
 - If symptoms are not severe, anaesthetists may be able to work in the clinical area.
- Areas where powdered gloves are used should be avoided.
- Neoprene/nitrile gloves should be used where there is a risk of blood contamination.
- Non-powdered latex or synthetic gloves should be worn by other staff in the area.
- Outside the work environment care also needs to be taken, particularly avoiding latex balloons and latex condoms!
- Be prepared if you become a patient !!

GUIDELINES FOR REDUCTION OF SENSITISATION TO LATEX

- Ongoing education regarding latex sensitivity and the safe use of gloves.
- Identification and advice to individuals with rashes or atopic conditions.
- Accurate diagnosis of reactions to latex gloves. It is important to see an allergist who has knowledge of this condition.
- Provision of appropriate gloves, ie powder-less low allergen latex gloves.

Thanks to Dr Helen Kolowale for this document

Further reading

Crippa M et al. Prevention of Latex allergy among health care workers: evaluation of the extractable latex protein content in different types of medical gloves. *Am J Ind Med* 2003; 44:24-31.

Latex allergy. A position paper of the British Society of Allergy and Clinical Immunology. *Clin Exp Allergy* 2003; 33:1484-1499.

This Resource Document has been prepared in good faith and having regard to general circumstances and is intended for information only. It is entirely the responsibility of the practitioner as to the manner in which s/he follows this document, having express regard to the circumstances of each case, and in the application of this document in each case.

The information contained in this document is not intended to constitute specific medical or other professional advice. The College and Societies, their officers, and employees, take no responsibility in relation to the application of use of this Resource Document in any particular circumstance.



The Resource Documents have been prepared having regard to the information available at the time of their preparation. They are reviewed from time to time, and it is the responsibility of the practitioner to ensure that s/he has obtained the current version. The practitioner should therefore have regard to any information, research or material which may have been published or become available subsequently.

Whilst the Welfare of Anaesthetists Special Interest Group endeavours to ensure that Resource Documents are as current as possible at the time of their preparation, it takes no responsibility for matters arising from changed circumstances or information or material which may have become available subsequently.

Promulgated: 1996
Date of current document: 2011

© This document is copyright; if it is reproduced in whole or in part, due acknowledgement is to be given.

ACECC is a joint initiative of the Australian and New Zealand College of Anaesthetists,
the Australian Society of Anaesthetists and the New Zealand Society of Anaesthetists.

ANZCA HOUSE 630 ST KILDA ROAD MELBOURNE VIC 3004
Telephone: (03) 9510 6299 Facsimile: (03) 9510 6786